# **TNPO1** Rabbit pAb

Catalog No.: A9435



## **Basic Information**

#### **Observed MW**

85-102kDa

#### **Calculated MW**

102kDa

#### Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes the beta subunit of the karyopherin receptor complex which interacts with nuclear localization signals to target nuclear proteins to the nucleus. The karyopherin receptor complex is a heterodimer of an alpha subunit which recognizes the nuclear localization signal and a beta subunit which docks the complex at nucleoporins. Alternate splicing of this gene results in several transcript variants encoding different proteins.

## **Recommended Dilutions**

**WB** 1:200 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**3842

Swiss Prot
Q92973

#### **Immunogen**

 $Synthetic\ peptide.\ This\ information\ is\ considered\ to\ be\ commercially\ sensitive.$ 

### **Synonyms**

MIP; TRN; IPO2; MIP1; KPNB2; TNPO1

## **Contact**

|          | 400-999-6126              |
|----------|---------------------------|
| <b>×</b> | cn.market@abclonal.com.cn |
|          | www.abclonal.com.cn       |

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using TNPO1 Rabbit pAb (A9435) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 1s.