# **ESRRG Rabbit mAb**

Catalog No.: A9606 Recombinant 1 Publications



## **Basic Information**

### **Observed MW**

47kDa

#### **Calculated MW**

51kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Mouse, Rat

### CloneNo number

ARC1653

# **Background**

This gene encodes a member of the estrogen receptor-related receptor (ESRR) family, which belongs to the nuclear hormone receptor superfamily. All members of the ESRR family share an almost identical DNA binding domain, which is composed of two C4-type zinc finger motifs. The ESRR members are orphan nuclear receptors; they bind to the estrogen response element and steroidogenic factor 1 response element, and activate genes controlled by both response elements in the absence of any ligands. The ESRR family is closely related to the estrogen receptor (ER) family. They share target genes, co-regulators and promoters, and by targeting the same set of genes, the ESRRs seem to interfere with the ER-mediated estrogen response in various ways. It has been reported that the family member encoded by this gene functions as a transcriptional activator of DNA cytosine-5methyltransferases 1 (Dnmt1) expression by direct binding to its response elements in the DNMT1 promoters, modulates cell proliferation and estrogen signaling in breast cancer, and negatively regulates bone morphogenetic protein 2-induced osteoblast differentiation and bone formation. Multiple alternatively spliced transcript variants have been identified, which mainly differ at the 5' end and some of which encode protein isoforms differing in the Nterminal region.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# Immunogen Information

Gene ID 2104 **Swiss Prot** 

P62508

### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human ESRRG (P62508).

## **Synonyms**

ERR3; ERRg; NR3B3; ERRgamma; ERR-gamma; ESRRG

# **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using ESRRG Rabbit mAb (A9606) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.