Collagen VI/COL6A1 Rabbit mAb

www.abclonal.com

ABclonal

Catalog No.: A9738 Recombinant 1 Publications

Basic Information

Observed MW

140kDa

Calculated MW

109kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC1725

Background

The collagens are a superfamily of proteins that play a role in maintaining the integrity of various tissues. Collagens are extracellular matrix proteins and have a triple-helical domain as their common structural element. Collagen VI is a major structural component of microfibrils. The basic structural unit of collagen VI is a heterotrimer of the alpha1(VI), alpha2(VI), and alpha3(VI) chains. The alpha2(VI) and alpha3(VI) chains are encoded by the COL6A2 and COL6A3 genes, respectively. The protein encoded by this gene is the alpha 1subunit of type VI collagen (alpha1(VI) chain). Mutations in the genes that code for the collagen VI subunits result in the autosomal dominant disorder, Bethlem myopathy.

Recommended Dilutions

WB 1:500 - 1:1000

1:50 - 1:200 **IHC-P**

Recommended starting **ELISA** concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 1291 P12109

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

OPLL; BTHLM1; UCHMD1; Collagen VI/COL6A1

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Product Information

Source Isotype **Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Western blot analysis of various lysates using Collagen VI/COL6A1 Rabbit mAb (A9738) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 3min.



Immunohistochemistry analysis of paraffinembedded Rat lung using Collagen VI/COL6A1 Rabbit mAb (A9738) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human liver using Collagen VI/COL6A1 Rabbit mAb (A9738) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.