# **HES5 Rabbit mAb**

Catalog No.: A9768 Recombinant 1 Publications



## **Basic Information**

### **Observed MW**

18kDa

### **Calculated MW**

18kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Mouse, Rat

#### CloneNo number

ARC1740

## **Background**

This gene encodes a member of a family of basic helix-loop-helix transcriptional repressors. The protein product of this gene, which is activated downstream of the Notch pathway, regulates cell differentiation in multiple tissues. Disruptions in the normal expression of this gene have been associated with developmental diseases and cancer.

# **Recommended Dilutions**

**WB** 1:1000 - 1:6000

IHC-P 1:100 - 1:500

**ELISA** Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**388585

Swiss Prot
Q5TA89

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

bHLHb38; HES5

## **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{a}$	ı	www.ahclonal.com.cn


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.




Western blot analysis of various lysates using HES5 Rabbit mAb (A9768) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunohistochemistry analysis of paraffinembedded Rat testis tissue using HES5 Rabbit mAb (A9768) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.