

β -Actin Rabbit mAb

Catalog No.: AC038

Recombinant

223 Publications

Basic Information

Observed MW

42kDa

Calculated MW

42kDa

Category

Loading control antibody

Applications

WB,IHC-P,ELISA

Cross-ReactivityHuman, Mouse, Rat, Chicken, Zebrafish,
Pig**CloneNo number**

ARC5115-01

Background

This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, integrity, and intercellular signaling. The encoded protein is a major constituent of the contractile apparatus and one of the two nonmuscle cytoskeletal actins that are ubiquitously expressed. Mutations in this gene cause Baraitser-Winter syndrome 1, which is characterized by intellectual disability with a distinctive facial appearance in human patients. Numerous pseudogenes of this gene have been identified throughout the human genome.

Recommended Dilutions

WB 1:10000 - 1:100000**IHC-P** 1:500 - 1:5000**ELISA** Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

60

Swiss Prot

P60709

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

SynonymsBRWS1; PS1TP5BP1; β -Actin

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification




Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Validation Data



Western blot analysis of lysates from HeLa cells, using β -Actin Rabbit mAb (AC038) at 1:10000-1:320000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using β -Actin Rabbit mAb (AC038) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using β -Actin Rabbit mAb (AC038) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using β -Actin Rabbit mAb (AC038) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat spleen tissue using β -Actin Rabbit mAb (AC038) at a dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.