

# Mouse anti DDDDK-Tag mAb

Catalog No.: AE005

281 Publications

## Basic Information

### Observed MW

37kDa/48kDa/57kDa/70kDa

### Calculated MW

### Category

Tag antibody

### Applications

WB,IF/ICC,IP,FC (intra),ELISA

### Cross-Reactivity

Species independent

### CloneNo number

AMC0382

## Background

FLAG-tag, or FLAG octapeptide, or FLAG epitope, is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. It has been used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits, because its mild purification procedure tends not to disrupt such complexes. It has been used to obtain proteins of sufficient purity and quality to carry out 3D structure determination by x-ray crystallography. A FLAG-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against a given protein, adding a FLAG-tag to a protein allows the protein to be studied with an antibody against the FLAG sequence. Examples are cellular localization studies by immunofluorescence or detection by SDS PAGE protein electrophoresis and Western blotting.

## Recommended Dilutions

**WB** 1:10000 - 1:80000**IF/ICC** 1:200 - 1:800**IP** 0.5µg-4µg antibody for  
100µg-200µg extracts of  
whole cells**FC (intra)** 5 µl per 10<sup>6</sup> cells in  
100 µl volume**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Immunogen Information

### Gene ID

### Swiss Prot

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

DDDDK;DDDDK tag;DDDDK-tag

## Product Information

### Source

Mouse

### Isotype

IgG1,Kappa

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.




[www.abclonal.com.cn](http://www.abclonal.com.cn)

---


# Validation Data



Immunoprecipitation of SERPINB1-Flag from 150 µg extracts of 293T cells transfected with a SERPINB1 expression vector containing a single N-terminal DDDDK-Tag was performed using 0.5 µg of Mouse anti DDDDK-Tag mAb (AE005). Mouse IgG isotype control (AC011) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10 % of the total input. Western blot analysis of immunoprecipitates was conducted using Mouse anti DDDDK-Tag mAb (AE005) at a dilution of 1:5000.





Western blot analysis of lysates from wild type (WT) and 293F cells transfected with Mouse anti DDDDK-Tag using Mouse anti DDDDK-Tag mAb (AE005) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates/proteins: 20 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45s.




Western blot analysis of lysates from wild type (WT) and 293F cells transfected with Mouse anti DDDDK-Tag using Mouse anti DDDDK-Tag mAb (AE005) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates/proteins: 20 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45s.


## Validation Data



Flow cytometry:  $1 \times 10^6$  CHO cells (negative control, left) and CHO-Claudin18.2-Flag (Transfection, right) cells were intracellularly-stained with Mouse anti DDDDK-Tag mAb (AE005, 2 µg/mL, orange line) or Mouse isotype control (2 µg/mL, blue line), followed by PE Goat anti-Mouse pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  CHO-Claudin18.2-Flag (Transfection, right) cells were intracellularly-stained with Mouse isotype control (2 µg/mL, left) or Mouse anti DDDDK-Tag mAb (AE005, 2 µg/mL, right), followed by PE Goat anti-Mouse pAb staining.



Confocal imaging of 293F cells transfected with DDDDK-Tag using Mouse anti DDDDK-Tag mAb (AE005, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L) (AS008, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.