

Mouse anti DDDDK-Tag mAb

Catalog No.: AE005 **249 Publications**

Basic Information

Observed MW

37kDa/48kDa/57kDa/70kDa

Calculated MW

Category

Tag antibody

Applications

WB,IF/ICC,IP,FC (intra),ELISA

Cross-Reactivity

Species independent

CloneNo number

AMC0382

Background

FLAG-tag, or FLAG octapeptide, or FLAG epitope, is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. It has been used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits, because its mild purification procedure tends not to disrupt such complexes. It has been used to obtain proteins of sufficient purity and quality to carry out 3D structure determination by x-ray crystallography. A FLAG-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against a given protein, adding a FLAG-tag to a protein allows the protein to be studied with an antibody against the FLAG sequence. Examples are cellular localization studies by immunofluorescence or detection by SDS PAGE protein electrophoresis and Western blotting.

Recommended Dilutions

WB 1:20000 - 1:80000

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
100µg-200µg extracts of
whole cells

FC (intra) 5 µl per 10⁶ cells in
100 µl volume

ELISA Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

Immunogen Information

Gene ID

Swiss Prot

Immunogen

A synthetic peptide corresponding to DDDDK tag.

Synonyms

DDDDK;DDDDK tag;DDDDK-tag

Product Information

Source

Mouse

Isotype

IgG1,Kappa

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.




| www.abclonal.com.cn


Validation Data



Immunoprecipitation of SERPINB1-Flag from 150 µg extracts of 293T cells transfected with a SERPINB1 expression vector containing a single N-terminal DDDDK-Tag was performed using 0.5 µg of Mouse anti DDDDK-Tag mAb (AE005). Mouse IgG isotype control (AC011) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10 % of the total input. Western blot analysis of immunoprecipitates was conducted using Mouse anti DDDDK-Tag mAb (AE005) at a dilution of 1:5000.




Western blot analysis of lysates from wild type (WT) and 293F cells transfected with Mouse anti DDDDK-Tag using Mouse anti DDDDK-Tag mAb (AE005) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates/proteins: 20 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45s.




Western blot analysis of lysates from wild type (WT) and 293F cells transfected with Mouse anti DDDDK-Tag using Mouse anti DDDDK-Tag mAb (AE005) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates/proteins: 20 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45s.


Validation Data




Western blot analysis of lysates from wild type (WT) and 293F cells transfected with Mouse anti DDDDK-Tag using Mouse anti DDDDK-Tag mAb (AE005) at 1:20000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L)(AS003) at 1:10000 dilution. Lysates/proteins: 20 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 45s.




Immunofluorescence analysis of 293T cells using Mouse anti DDDDK-Tag mAb (AE005) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunofluorescence analysis of 293T-BCAT2-Flag-GFP-C cells using Mouse anti DDDDK-Tag mAb (AE005) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of 293T-BCAT2-Flag-GFP-N cells using Mouse anti DDDDK-Tag mAb (AE005) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Flow cytometry: 1×10^6 CHO cells (negative control, left) and CHO-Claudin18.2-Flag (Transfection, right) cells were intracellularly-stained with Mouse anti DDDDK-Tag mAb (AE005, 2 µg/mL, orange line) or Mouse isotype control (2 µg/mL, blue line), followed by PE Goat anti-Mouse pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 CHO-Claudin18.2-Flag (Transfection, right) cells were intracellularly-stained with Mouse isotype control (2 µg/mL, left) or Mouse anti DDDDK-Tag mAb (AE005, 2 µg/mL, right), followed by PE Goat anti-Mouse pAb staining.