

Rabbit anti GST-Tag pAb

Catalog No.: AE006

36 Publications

Basic Information

Observed MW

27kDa

Calculated MW

26kDa

Category

Tag antibody

Applications

WB, IP

Cross-Reactivity

Species independent

Background

Glutathione S-transferases (GSTs), previously known as ligandins, comprise a family of eukaryotic and prokaryotic phase II metabolic isozymes best known for their ability to catalyze the conjugation of the reduced form of glutathione (GSH) to xenobiotic substrates for the purpose of detoxification. The GST family consists of three superfamilies: the cytosolic, mitochondrial, and microsomal—also known as MAPEG—proteins. Members of the GST superfamily are extremely diverse in amino acid sequence, and a large fraction of the sequences deposited in public databases are of unknown function. The Enzyme Function Initiative (EFI) is using GSTs as a model superfamily to identify new GST functions. A GST-tag is often used to separate and purify proteins that contain the GST-fusion protein. The tag is 220 amino acids (roughly 26 kDa) in size, which, compared to tags such as the Myc-tag or the FLAG-tag, is quite large. It can be fused to either the N-terminus or C-terminus of a protein. However, many commercially available sources of GST-tagged plasmids include a thrombin domain for cleavage of the GST tag during protein purification.

Recommended Dilutions

WB 1:1000 - 1:2000**IP** 3µg antibody for 50µg
extracts of recombinant
protein

Immunogen Information

Gene ID**Swiss Prot**

P08515

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

GST; GST tag; GST-tag

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Immunoprecipitation analysis of 50 µg extracts of GST Protein using 3 µg GST-Tag antibody (AE006). Western blot was performed from the immunoprecipitate using GST-Tag antibody (AE006).



Western blot analysis of over-expressed GST protein using Rabbit anti GST-Tag pAb (AE006) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Western blot analysis of over-expressed GST protein using Rabbit anti GST-Tag pAb (AE006) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.