

Mouse anti HA-Tag mAb

Catalog No.: AE008

267 Publications

Basic Information

Observed MW

26 kDa

Calculated MW

Category

Tag antibody

Applications

WB,IP,IF/ICC,ELISA

Cross-Reactivity

Species independent

CloneNo number

AMC0503

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB 1:10000 - 1:20000**IP** 0.5 µg-4 µg antibody for
200 µg-400 µg extracts
of whole cells**IF/ICC** 1:50 - 1:200**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

HA;HA tag;HA-tag

Product Information

Source

Mouse

Isotype

IgG1,Kappa

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of over-expressed HA-tagged protein in 293T cell using Mouse anti HA-Tag mAb (AE008) at different dilution. Each lane was loaded with 2 ug cell lysate.
 Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 1s.



Western blot analysis of lysates from 293T cells, using Mouse anti HA-Tag mAb (AE008) at 1:10000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 10s.



Immunofluorescence analysis of 293T cells transfected with HA-Tag fusion protein and untreated 293T cells use Mouse anti HA-Tag mAb (AE008) at dilution of 1:50 (40x lens).
 Blue: DAPI for nuclear staining.



Immunoprecipitation of over-expressed HA-tagged protein in 293T cells using HA-tag antibody (AE008). A mock served as negative control and over-expressed 293T cell lysate served as positive control.