Mouse anti Myc-Tag mAb

Catalog No.: AE010 94 Publications



Basic Information

Observed MW

50-70kDa

Calculated MW

Category

Tag antibody

Applications

ELISA,WB,IP

Cross-Reactivity

Species independent

CloneNo number

AMC0504

Background

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

Recommended Dilutions

WB 1:2000 - 1:6000

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

Immunogen Information

Gene ID Swiss Prot

Immunogen

A synthetic peptide corresponding to Myc tag.

Synonyms

Myc;Myc tag;Myc-tag

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\Box}$	www.ahclonal.com.cn

Product Information


SourceIsotypePurificationMouseIgG1Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of normal 293T cells and 293T transfected with Myc-NLK protein, using Mouse anti Myc-Tag mAb (AE010) at 1:5000 dilution.

Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.