

# Rabbit anti His-tag mAb

Catalog No.: AE086 27 Publications

## Basic Information

### Observed MW

35 kDa, 58 kDa

### Calculated MW

### Category

Tag antibody

### Applications

WB, IF/ICC, IP, ELISA

### Cross-Reactivity

Species independent

### CloneNo number

ARC2696

## Background

Consecutive histidine residues (usually 6 to 10 in length) are often inserted into the amino acid sequences of recombinant proteins. The resulting His-tagged proteins can be detected by using His antibodies.

## Recommended Dilutions

WB	1:10000 - 1:60000
IF/ICC	1:50 - 1:200
IP	0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>

## Immunogen Information

### Gene ID

### Swiss Prot

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

His;His tag;His-tag

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data



Immunoprecipitation analysis of 300 µg extracts of 293T cells using 3 µg His-tag antibody (AE086). Western blot was performed from the immunoprecipitate using His-tag antibody (AE086) at a dilution of 1:30000.



Western blot analysis of lysates from wild type (WT) 293F cells, 293F cells transfected with Snail-His-tag and 293F cells transfected with CYP1A2-His-tag using Rabbit anti His-tag mAb (AE086) at 1:11000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 20 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020)

Exposure time: 10 s.



Immunofluorescence analysis of 293T-SOST His(C) and 293T cells using Rabbit anti His-tag mAb (AE086) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.