# Phospho-EGFR-Y1148 Rabbit pAb

Catalog No.: AP0028



## **Basic Information**

#### **Observed MW**

180kDa

#### **Calculated MW**

134kDa

#### Category

Primary antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

## **Background**

The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor, thus inducing receptor dimerization and tyrosine autophosphorylation leading to cell proliferation. Mutations in this gene are associated with lung cancer. EGFR is a component of the cytokine storm which contributes to a severe form of Coronavirus Disease 2019 (COVID-19) resulting from infection with severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2).

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
1956
P00533

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

ERBB; ERRP; HER1; mENA; ERBB1; PIG61; NISBD2; Phospho-EGFR-Y1148

## **Contact**

<b>a</b>	400-999-6126
$\sim$	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

# **Validation Data**



Western blot analysis of lysates from A-431 cells, using Phospho-EGFR-Y1148 Rabbit pAb (AP0028) at 1:1000 dilution. A431 cells were treated with EGF (100 ng/mL) for 30 minutes after serum-starvation overnight.

 $Secondary\ antibody:\ HRP\text{-}conjugated\ Goat\ anti-Rabbit\ IgG\ (H+L)\ (AS014)\ at\ 1:10000\ dilution.$ 

Lysates/proteins:  $25\mu g$  per lane.

Blocking buffer: 3% BSA.