# Phospho-c-Jun-S63 Rabbit pAb

Catalog No.: AP0048 20 Publications



## **Basic Information**

#### **Observed MW**

48kDa

## **Calculated MW**

36kDa

## Category

Primary antibody

## **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Rat

# **Background**

This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. This gene is intronless and is mapped to 1p32p31, a chromosomal region involved in both translocations and deletions in human malignancies.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID Swiss Prot** 3725 P05412

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

AP1; p39; AP-1; cJUN; c-Jun; Phospho-c-Jun-S63

## **Contact**

2		400-999-6126
$\bowtie$	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

## **Product Information**


**Purification** Source Isotype Rabbit IgG Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

# **Validation Data**



Western blot analysis of lysates from C6 cells, using Phospho-c-Jun-S63 Rabbit pAb (AP0048) at 1:1000 dilution. C6 cells were treated with UV at room temperature for 15-30 minutes.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.