# Phospho-PRKCQ-S695 Rabbit pAb

Catalog No.: AP0192 1 Publications



## **Basic Information**

#### **Observed MW**

72kDa

#### **Calculated MW**

82kDa

#### Category

Primary antibody

## **Applications**

WB

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase is important for T-cell activation. It is required for the activation of the transcription factors NF-kappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of the transcription factors.

## **Recommended Dilutions**

**WB** 

1:500 - 1:2000

## Immunogen Information

Gene ID 5588 Swiss Prot Q04759

#### **Immunogen**

A phospho specific peptide corresponding to residues surrounding S695 of human PRKCQ

## **Synonyms**

PRKCT; nPKC-theta; Phospho-PRKCQ-S695

## **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts from 293 cells, using Phospho-PRKCQ-S695 antibody (AP0192). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins:  $25\mu g$  per lane. Blocking buffer: 3% BSA.