Leader in Biomolecular Solutions for Life Science

# Phospho-c-Abl-Y412 Rabbit pAb

Catalog No.: AP0303



# **Basic Information**

**Observed MW** Refer to figures

Calculated MW 123kDa

Category Primary antibody

Applications IHC-P,ELISA

Cross-Reactivity Human, Mouse, Rat

# Background

This gene is a protooncogene that encodes a protein tyrosine kinase involved in a variety of cellular processes, including cell division, adhesion, differentiation, and response to stress. The activity of the protein is negatively regulated by its SH3 domain, whereby deletion of the region encoding this domain results in an oncogene. The ubiquitously expressed protein has DNA-binding activity that is regulated by CDC2-mediated phosphorylation, suggesting a cell cycle function. This gene has been found fused to a variety of translocation partner genes in various leukemias, most notably the t(9;22) translocation that results in a fusion with the 5' end of the breakpoint cluster region gene (BCR; MIM:151410). Alternative splicing of this gene results in two transcript variants, which contain alternative first exons that are spliced to the remaining common exons.

# **Recommended Dilutions**

IHC-P	1:50 - 1:100
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

Gene ID

Swiss Prot P00519

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms** 

ABL; JTK7; p150; c-ABL; v-abl; CHDSKM; c-ABL1; BCR-ABL; bcr/abl; Phospho-c-Abl-Y412

# Contact


6	400-999-6126
$\times$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

# **Product Information**

**Source** Rabbit **lsotype** IgG Purification Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Immunohistochemistry analysis of paraffinembedded Human breast carcinoma using Phospho-c-Abl-Y412 Rabbit pAb (AP0303). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.