

# Phospho-CDKN1B/p27KIP1-T187 Rabbit pAb

Catalog No.: AP0328

## Basic Information

### Observed MW

26kDa

### Calculated MW

22kDa

### Category

Primary antibody

### Applications

WB

### Cross-Reactivity

Human

## Background

This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. Mutations in this gene are associated with multiple endocrine neoplasia type IV (MEN4).

## Recommended Dilutions

WB 1:500 - 1:1000

## Immunogen Information

### Gene ID

1027

### Swiss Prot

P46527

### Immunogen

A phospho specific peptide corresponding to residues surrounding T187 of human CDKN1B/p27KIP1

### Synonyms

KIP1; MEN4; CDKN4; MEN1B; P27KIP1; Phospho-CDKN1B/p27KIP1-T187

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of lysates from HeLa cells, using Phospho-CDKN1B/p27KIP1-T187 Rabbit pAb (AP0328).  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% BSA.