

# Phospho-eIF2 $\alpha$ -S49 Rabbit pAb

Catalog No.: AP0341 **5 Publications**

## Basic Information

**Observed MW**

40kDa

**Calculated MW**

36kDa

**Category**

Primary antibody

**Applications**

WB, IHC-P

**Cross-Reactivity**

Human, Mouse, Rat

## Background

The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).

## Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:100

## Immunogen Information

**Gene ID**

1965

**Swiss Prot**

P05198

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**EIF2; EIF-2; EIF2A; EIF-2A; EIF-2alpha; Phospho-eIF2 $\alpha$ -S49

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data



Western blot analysis of lysates from HeLa cells, using Phospho-eIF2 $\alpha$ -S49 Rabbit pAb (AP0341).  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% BSA.



Immunohistochemistry analysis of paraffin-embedded Human breast carcinoma using Phospho-eIF2 $\alpha$ -S49 Rabbit pAb (AP0341).  
Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.