

Phospho-ZAP70-Y319 Rabbit pAb

Catalog No.: AP0467

Basic Information

Observed MW

70kDa

Calculated MW

70kDa

Category

Primary antibody

Applications

ELISA, WB, IP

Cross-Reactivity

Human

Background

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID

7535

Swiss Prot

P43403

Immunogen

A synthetic phosphorylated peptide around Y319 of human ZAP70 (NP_001070.2).

Synonyms

SRK; STD; TZK; STCD; IMD48; ADMIO2; ZAP-70; Phospho-ZAP70-Y319

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data



Immunoprecipitation analysis of 200 μ g extracts of Jurkat cells, using 3 μ g Phospho-ZAP70-Y319 pAb (AP0467). Western blot was performed from the immunoprecipitate using Phospho-ZAP70-Y319 pAb (AP0467) at a dilution of 1:1000. Jurkat cells were treated by Hydrogen Peroxide (2 mM) at 37°C for 2 minutes after serum-starvation overnight.