# Phospho-AURKA-T288 Rabbit pAb

Catalog No.: AP0523 2 Publications



## **Basic Information**

#### **Observed MW**

48kDa

#### **Calculated MW**

46kDa

### Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript variants encoding the same protein have been found for this gene.

# **Recommended Dilutions**

**WB** 

1:500 - 1:2000

# **Immunogen Information**

Gene ID 6790 **Swiss Prot** 

014965

#### **Immunogen**

A synthetic phosphorylated peptide around T288 of human AURKA (NP $\_$ 003591.2).

## **Synonyms**

AIK; ARK1; AURA; BTAK; STK6; STK7; STK15; PPP1R47; Phospho-AURKA-T288

# **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates, using Phospho-AURKA-T288 Rabbit pAb (AP0523) at 1:1000 dilution. HeLa were treated by Hydroxyurea (4mM) or treated by Nocodazole (50ng/mL) for 20 hours. HT-29 cells were treated by Hydroxyurea (4 mM) for 20 hours or treated by Nocodazole (100ng/mL) for 16 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.