

Phospho-TrkA-Y674/675 + TrkB-Y706/707 Rabbit pAb

Catalog No.: AP0549

Basic Information

Observed MW

140kDa

Calculated MW

35kDa/53-60kDa/77kDa/83-93kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human

Background

This gene encodes a member of the neurotrophic tyrosine kinase receptor (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date. [provided by RefSeq, Jul 2008]

Recommended Dilutions

WB 1:100 - 1:500

Immunogen Information

Gene ID

4914/4915

Swiss Prot

P04629/Q16620

Immunogen

A synthetic phosphorylated peptide around Y674 & Y675 of human TrkANTRK1 (NP_001012331.1).

Synonyms

NTRK1/NTRK2; Phospho-TrkA-Y674/675 + TrkB-Y706/707

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from HeLa cells, using Phospho-TrkA-Y674/675 + TrkB-Y706/707 Rabbit pAb (AP0549) at 1:500 dilution. HeLa cells were treated by Pervanadate (1 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.