Phospho-ALOX5-S271 Rabbit pAb

Catalog No.: AP0573



Basic Information

Observed MW

78kDa

Calculated MW

78kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human

Background

This gene encodes a member of the lipoxygenase gene family and plays a dual role in the synthesis of leukotrienes from arachidonic acid. The encoded protein, which is expressed specifically in bone marrow-derived cells, catalyzes the conversion of arachidonic acid to 5(S)-hydroperoxy-6-trans-8,11,14-cis-eicosatetraenoic acid, and further to the allylic epoxide 5(S)-trans-7,9-trans-11,14-cis-eicosatetrenoic acid (leukotriene A4). Leukotrienes are important mediators of a number of inflammatory and allergic conditions. Mutations in the promoter region of this gene lead to a diminished response to antileukotriene drugs used in the treatment of asthma and may also be associated with atherosclerosis and several cancers. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID240

Swiss Prot
P09917

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

5-LO; 5LPG; LOG5; 5-LOX; Phospho-ALOX5-S271

Contact

<u>a</u>	400-999-6126
×	cn.market@abclonal.com.cn
$\overline{\mathfrak{S}}$	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from HT-29 cells, using Phospho-ALOX5-S271 Rabbit pAb (AP0573) at 1:1000 dilution. HT-29 cells were treated by PMA/TPA (200nM).

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.

Antibody | Protein | ELISA Kits | Enzyme | NGS | Service