# Phospho-eIF4E-S209 Rabbit pAb

Catalog No.: AP0747 1 Publications



# **Basic Information**

#### **Observed MW**

25kDa

#### **Calculated MW**

25kDa

## Category

Primary antibody

#### **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse

# **Background**

The protein encoded by this gene is a component of the eukaryotic translation initiation factor 4F complex, which recognizes the 7-methylguanosine cap structure at the 5' end of messenger RNAs. The encoded protein aids in translation initiation by recruiting ribosomes to the 5'-cap structure. Association of this protein with the 4F complex is the rate-limiting step in translation initiation. This gene acts as a proto-oncogene, and its expression and activation is associated with transformation and tumorigenesis. Several pseudogenes of this gene are found on other chromosomes. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

**WB** 

1:100 - 1:500

# Immunogen Information

Gene ID 1977 Swiss Prot

P06730

# Immunogen

A synthetic phosphorylated peptide around S209 of human eIF4E (NP\_001959.1).

#### **Synonyms**

CBP; EIF4F; AUTS19; EIF4E1; eIF-4E; EIF4EL1; Phospho-eIF4E-S209

### **Contact**

<b>a</b>	400-999-6126
$\sim$	cn.market@abclonal.com.cn
<b>⊙</b>	www.abclonal.com.cn

#### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-elF4E-S209 Rabbit pAb (AP0747) at 1:500 dilution. NIH/3T3 cells were treated by 10% FBS at  $37^{\circ}$ C for 30 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 3min.