

# Phospho-Glucocorticoid Receptor-S211 Rabbit pAb

Catalog No.: AP0759 **1 Publications**

## Basic Information

**Observed MW**

95kDa

**Calculated MW**

86kDa

**Category**

Primary antibody

**Applications**

ELISA,WB

**Cross-Reactivity**

Human,Mouse

## Background

This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175).

## Recommended Dilutions

WB 1:1000 - 1:2000

## Immunogen Information

**Gene ID**

2908

**Swiss Prot**

P04150

**Immunogen**

A synthetic phosphorylated peptide around S211 of human Glucocorticoid Receptor (NP\_000167.1).

**Synonyms**

GR; GCR; GRL; GCCR; GCRST; Phospho-Glucocorticoid Receptor-S211

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**

Affinity purification


**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data

---



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-Glucocorticoid Receptor-S211 Rabbit pAb (AP0759) at 1:600 dilution. NIH/3T3 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 30s.