

Phospho-PI3KP85 α /P55 γ /P85 β -Y467/Y199/Y464 Rabbit pAb

Catalog No.: AP0854

82 Publications

Basic Information

Observed MW

60kDa(P55)/85kDa(P85)

Calculated MW

42-54kDa/81-84kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit.

Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in four transcript variants encoding different isoforms. [provided by RefSeq, Jun 2011]

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

5295/5296/8503

Swiss Prot

P27986/O00459/Q92569

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PIK3R1/PIK3R2/PIK3R3; Phospho-PI3KP85 α /P55 γ /P85 β -Y467/Y199/Y464

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from wild type (WT) and 293T cells were transfected with SRC(Y530F), using Phospho-PI3KP85α/γ/β-Y467/Y199/Y464 Rabbit pAb (AP0854) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Western blot analysis lysates from C2C12, using Phospho-PI3KP85α/P55γ/P85β-Y467/Y199/Y464 Rabbit pAb (AP0854) at 1:1000 dilution. C2C12 cells were treated with Hydrogen Peroxide (2 nM) at 37°C for 15 minutes after serum-starvation overnight. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.