

Phospho-PERK-T982 Rabbit pAb

Catalog No.: AP0886

39 Publications

Basic Information

Observed MW

170kDa/

Calculated MW

125kDa

Category

Primary antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene phosphorylates the alpha subunit of eukaryotic translation-initiation factor 2, leading to its inactivation, and thus to a rapid reduction of translational initiation and repression of global protein synthesis. This protein is thought to modulate mitochondrial function. It is a type I membrane protein located in the endoplasmic reticulum (ER), where it is induced by ER stress caused by malformed proteins. Mutations in this gene are associated with Wolcott-Rallison syndrome.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

9451

Swiss Prot

Q9NZJ5

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PEK; WRS; PERK; Phospho-PERK-T982

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from A549 cells using Phospho-PERK-T982 Rabbit pAb (AP0886) at 1:1000 dilution incubated overnight at 4°C. A549 cells were treated with 10% FBS at 37°C for 5 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Western blot analysis of lysates from 293T cells, using Phospho-PERK-T982 Rabbit pAb (AP0886) at 1:1000 dilution. 293T cells were treated with IGF-1 (50 ng/ml) at 37°C for 5 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.




Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using Phospho-PERK-T982 Rabbit pAb (AP0886) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat lung tissue using Phospho-PERK-T982 Rabbit pAb (AP0886) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse Intestine tissue using Phospho-PERK-T982 Rabbit pAb (AP0886) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using Phospho-PERK-T982 Rabbit pAb (AP0886) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.