ABclonal www.abclonal.com

Pan Phospho-Tyrosine Rabbit pAb

Catalog No.: AP0905 6 Publications

Basic Information

Observed MW

20-70kDa

Calculated MW

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Tyrosine phosphorylation (pTyr), much of which occurred on localized multiple sites, initiates cellular signaling, governs cellular functions, and its dysregulation is implicated in many diseases, especially cancers. pTyr-specific sensing is of great significance for understanding disease states and developing targeted anticancer drugs.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

 $\begin{array}{c} \textbf{ELISA} & \text{Recommended starting} \\ & \text{concentration is 1 } \mu\text{g/mL}. \end{array}$

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\overline{a}	ī	www.ahclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 50% glycerol,pH7.3.




Western blot analysis of lysates from Jurkat cells, using Pan Phospho-Tyrosine pAb (AP0905) at 1:1000 dilution. Jurkat cells were treated with Pervanadate (1 mM) at 37° C for 30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of C6 cells using Pan Phospho-Tyrosine (AP0905) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.C6 cells were treated with 20% FBS at 37 $^{\circ}$ C for 20 minutes after serum-starvation overnight.



Immunofluorescence analysis of NIH/3T3 cells using Pan Phospho-Tyrosine (AP0905) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.NIH/3T3 cells were treated with 20% FBS at 37°C for 20 minutes after serum-starvation overnight.