Phospho-PKC-beta-T641 Rabbit pAb

Catalog No.: AP0971



Basic Information

Observed MW

Refer to figures

Calculated MW

77kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively spliced transcript variants encoding distinct isoforms have been reported.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID5579 **Swiss Prot**P05771

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PKCB; PRKCB1; PRKCB2; PKCI(2); PKCbeta; PKC-beta; Phospho-PKC-beta-T641

Contact

| a | 400-999-6126 |
|-------------------|---------------------------|
| \sim | cn.market@abclonal.com.cn |
| $\overline{\Box}$ | www.ahclonal.com.cn |

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.