

# Phospho-Tyrosine Hydroxylase-S40 Rabbit pAb

Catalog No.: AP1109

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

59kDa

### Category

Primary antibody

### Applications

ELISA, WB

### Cross-Reactivity

Rat

## Background

The protein encoded by this gene is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in this gene have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.

## Recommended Dilutions

WB 1:500 - 1:2000

## Immunogen Information

### Gene ID

7054

### Swiss Prot

P07101

### Immunogen

A synthetic phosphorylated peptide around S40 of human Tyrosine Hydroxylase (NP\_000351.2?).

### Synonyms

TYH; DYT14; DYT5b; Phospho-Tyrosine Hydroxylase-S40

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

## Validation Data

---



Western blot analysis of lysates from PC-12 cells, using Phospho-Tyrosine Hydroxylase-S40 pAb (AP1109) at 1:1000 dilution or Tyrosine Hydroxylase antibody (A0028). PC-12 cells were treated by Forskolin (30  $\mu$ M) at 37°C for 30 minutes.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.