Phospho-Vimentin-S83 Rabbit pAb

Catalog No.: AP1119



Basic Information

Observed MW

57kDa

Calculated MW

54kDa

Category

Primary antibody

Applications

WB,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a type III intermediate filament protein. Intermediate filaments, along with microtubules and actin microfilaments, make up the cytoskeleton. The encoded protein is responsible for maintaining cell shape and integrity of the cytoplasm, and stabilizing cytoskeletal interactions. This protein is involved in neuritogenesis and cholesterol transport and functions as an organizer of a number of other critical proteins involved in cell attachment, migration, and signaling. Bacterial and viral pathogens have been shown to attach to this protein on the host cell surface. Mutations in this gene are associated with congenital cataracts in human patients.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID7431

Swiss Prot
P08670

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

VIM; CTRCT30; HEL113; vimentin; Phospho-Vimentin-S83

Contact

| a | | 400-999-6126 |
|-----------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| \odot | T | www.abclonal.com.cn |


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from NIH/3T3 cells, using Phospho-Vimentin-S83 Rabbit pAb (AP1119) at 1:1000 dilution. NIH/3T3 cells were treated with Paclitaxel (100 nM/ml) at 37°C for 20 hours. NIH/3T3 cells were treated with Hydroxyurea (4 mM) at 37 $^{\circ}\text{C}$ for 20 hours.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of lysates from C6 cells, using Phospho-Vimentin-S83 Rabbit pAb (AP1119) at 1:1000 dilution. C6 cells were treated with Paclitaxel (100 nM) at 37°C for 20 hours.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Western blot analysis of lysates from HeLa cells, using Phospho-Vimentin-S83 Rabbit pAb (AP1119) at 1:1000 dilution. HeLa cells were treated with Paclitaxel (100 nM/ml) at 37°C for 20 hours. HeLa cells were treated with Hydroxyurea (4 mM) at 37°C for 20 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.