

Phospho-S6 Ribosomal Protein (RPS6)-S235/236 Rabbit mAb

Catalog No.: AP1325

Recombinant

Basic Information

Observed MW

32kDa

Calculated MW

29kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC53843

Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Recommended Dilutions

WB 1:2000 - 1:7000**IHC-P** 1:100 - 1:500

Immunogen Information

Gene ID

6194

Swiss Prot

P62753

Immunogen

A synthetic phosphorylated peptide around S235 & S236 of human S6 Ribosomal Protein (RPS6) (NP_001001.2).

Synonyms

S6; eS6; Phospho-S6 Ribosomal Protein (RPS6)-S235/236

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data




Western blot analysis of extracts of NIH/3T3, using Phospho-S6 Ribosomal Protein (RPS6)-S235/236 antibody (AP1325) at 1:6000 dilution. NIH/3T3 cells were treated by CIP (20 uL/400 uL) at 37°C for 1 hour. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Western blot analysis of extracts of PC-12, using Phospho-S6 Ribosomal Protein (RPS6)-S235/236 antibody (AP1325) at 1:6000 dilution. PC-12 cells were treated by 10% FBS at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Immunohistochemistry analysis of paraffin-embedded human liver cancer using Phospho-S6 Ribosomal Protein (RPS6)-S235/236 Rabbit mAb (AP1325) at dilution of 1:500 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using Phospho-S6 Ribosomal Protein (RPS6)-S235/236 Rabbit mAb (AP1325) at dilution of 1:500 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.