

Phospho-S6 Ribosomal Protein (RPS6)-S240/244 Rabbit mAb

Catalog No.: AP1328 **Recombinant** **4 Publications**

Basic Information

Observed MW

32kDa

Calculated MW

29kDa

Category

Primary antibody

Applications

WB,IF/ICC,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC54426

Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Recommended Dilutions

WB 1:20000 - 1:200000

IF/ICC 1:200 - 1:1000

IHC-P 1:100 - 1:500

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

6194

Swiss Prot

P62753

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

S6; eS6; Phospho-S6 Ribosomal Protein (RPS6)-S240/244

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Western blot analysis of various lysates using Phospho-S6 Ribosomal Protein (RPS6)-S240/244 Rabbit mAb (AP1328) at 1:20000 dilution. HeLa and NIH/3T3 cell were treated with 10% FBS at 37°C for 30 minutes after serum-starvation overnight. PC-12 cells were treated with 20% FBS at 37°C for 20 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using Phospho-S6 Ribosomal Protein (RPS6)-S240/244 Rabbit mAb (AP1328) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Confocal imaging of PC-12 cells (treated with 20% FBS) and PC-12 cells (untreated) using Phospho-S6 Ribosomal Protein (RPS6)-S240/244 Rabbit mAb (AP1328, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.