

# Phospho-Jak1-Y1034/1035 Rabbit mAb

Catalog No.: AP1399

Recombinant

1 Publications

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

133kDa

### Category

Primary antibody

### Applications

WB,(ELISA)

### Cross-Reactivity

Human

### CloneNo number

ARC3147

## Background

This gene encodes a membrane protein that is a member of a class of protein-tyrosine kinases (PTK) characterized by the presence of a second phosphotransferase-related domain immediately N-terminal to the PTK domain. The encoded kinase phosphorylates STAT proteins (signal transducers and activators of transcription) and plays a key role in interferon-alpha/beta, interferon-gamma, and cytokine signal transduction. This gene plays a crucial role in effecting the expression of genes that mediate inflammation, epithelial remodeling, and metastatic cancer progression. This gene is a key component of the interleukin-6 (IL-6)/JAK1/STAT3 immune and inflammation response and is a therapeutic target for alleviating cytokine storms. The kinase activity of this gene is directly inhibited by the suppressor of cytokine signalling 1 (SOCS1) protein. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

3716

### Swiss Prot

P23458

### Immunogen

A synthesized peptide derived from human Jak1.

### Synonyms

JTK3; AIIDE; JAK1A; JAK1B; Phospho-Jak1-Y1022/1023

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Protein A purified


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PSB with 0.01% Sodium azide,0.05% BSA□40% glycerol,pH7.2.

## Validation Data

---



Western blot analysis of lysates from Jurkat cells, using Phospho-Jak1-Y1034/1035 Rabbit mAb (AP1399) at 1:1000 dilution. Jurkat cells were treated by Pervanadate (1 mM) at 37°C for 30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.