

Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb

Catalog No.: AP1430 **Recombinant**

Basic Information

Observed MW

60kDa

Calculated MW

48kDa/51kDa/55kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC61098

Background

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene.

Recommended Dilutions

WB	1:2000 - 1:10000
IHC-P	1:100 - 1:500

Immunogen Information

Gene ID

207/208/10000

Swiss Prot

P31749/P31751/Q9Y243

Immunogen

A synthetic phosphorylated peptide around T450 of human Akt.

Synonyms

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from RD cells, using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at 1:10000 dilution. RD cells were treated by λ -PP mixed solution (1ul) at 30°C for 30 minutes.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of lysates from C2C12 cells, using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at 1:10000 dilution. C2C12 cells were treated by λ -PP mixed solution (1ul) at 30°C for 30 minutes.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.



Immunohistochemistry analysis of Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 in paraffin-embedded Human breast tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 in paraffin-embedded Human tonsil tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 in paraffin-embedded Mouse intestine tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


Validation Data



Immunohistochemistry analysis of Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 in paraffin-embedded Mouse kidney tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 in paraffin-embedded Rat intestine tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 in paraffin-embedded Rat kidney tissue using Phospho-AKT1-T450 + AKT2-T451 + AKT3-T447 Rabbit mAb (AP1430) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.