

Phospho-AMPK α 1-T183+AMPK α 2-T172 Rabbit mAb

Catalog No.: AP1441 **Recombinant**

Basic Information

Observed MW

62kDa

Calculated MW

64kDa/65kDa/62kDa

Category

Primary antibody

Applications

ELISA, WB

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC62278

Background

The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

Recommended Dilutions

WB 1:1000 - 1:5000

Immunogen Information

Gene ID

5562/5563

Swiss Prot

Q13131/P54646

Immunogen

A synthetic phosphorylated peptide around T183/T172 of human PRKAA1/PRKAA2 (NP_006242.5/NP_006243.2).

SynonymsAMPK α 1/AMPK α 2; Phospho-AMPK α 1-T183+AMPK α 2-T172

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from C2C12 cells, using Phospho-AMPKα1-T183/AMPKα2-T172 Rabbit mAb (AP1441) at 1:1000 dilution. C2C12 cells were treated by mHGF (50 ng/uL).
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Western blot analysis of lysates from C6 cells, using Phospho-AMPKα1-T183/AMPKα2-T172 Rabbit mAb (AP1441) at 1:1000 dilution. C6 cells were treated by AICAR (0.5 mM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.