

ABflo® 647-conjugated Goat anti-Mouse IgG (H+L)

Catalog No.: AS059 **2 Publications**

Basic Information

Observed MW

Calculated MW

Category

Secondary antibody

Applications

IF/ICC,FC

Cross-Reactivity

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies. Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

Recommended Dilutions

IF/ICC 1:100 - 1:500

FC 1:50 - 1:100

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Mouse IgG

Synonyms

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Goat

Isotype

IgG

Purification




Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,1% BSA,50% glycerol,pH7.3.

Validation Data



Flow cytometric analysis of Positive antibody Human Calcitonin R (2.5µg/mL) in various cells (orange) compare to Mouse isotype control (blue) and non-staining control (Red). The secondary antibody used was ABflo® 647-conjugated Goat Anti-Mouse IgG (H+L) (AS059) at 1:100.

Confocal imaging of HeLa cells using α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by a further incubation with ABflo® 647-conjugated Goat Anti-Mouse IgG (H+L) (AS059, dilution 1:200) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.