

# HRP-conjugated Rabbit anti-Human IgG1 (Fc) mAb

Catalog No.: AS092

### **Basic Information**

#### **Observed MW**

50 kDa

#### **Calculated MW**

### Category

Secondary antibody

### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human

#### CloneNo number

ARC53385-HRP

### Conjugate

 $\mathsf{HRP}$ 

# **Background**

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies . Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

### **Recommended Dilutions**

**WB** 1:2000 - 1:30000

**ELISA** 1:5000-1:10000

### Immunogen Information

Gene ID Swiss Prot

#### **Immunogen**

This information is considered to be commercially sensitive.

#### **Synonyms**

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Т	www.abclonal.com.cn

### **Product Information**


Source Isotype Purification
Rabbit Horseradish peroxidase conjugated IgG

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3

## **Validation Data**



Western blot analysis of vlysates from Human plasma using HRP-conjugated Rabbit anti-Human IgG1 (Fc) mAb (AS092) at 1:4000-1:32000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30 s.