Leader in Biomolecular Solutions for Life Science

www.abclonal.com

APC Streptavidin

Catalog No.: AT0008

Basic Information

Observed MW

Calculated MW

Category
Primary antibody

Applications
FC

Cross-Reactivity
Species independent

CloneNo number
Streptavidin

Conjugate
APC. Ex:650nm. Em:660nm.

Background

Streptavidin binds to biotin with high affinity. Streptavidin-APC is useful for detecting biotinylated antibodies.

Recommended Dilutions

FC	≤0.25 μg per million cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen
This information is considered to be commercially sensitive.

Synonyms
SA V1; SA V2

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype


Purification
Affinity purification

Storage
Store at 2-8℃

Validation Data



Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with Biotin Rabbit IgG isotype control (5 µl/Test,left) or Biotin Rabbit anti-Mouse CD8a mAb (A27373,5 µl/Test,right), followed by APC Streptavidin (AT0008,0.25 µg) staining.



Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with Biotin Rabbit anti-Mouse CD8a mAb (A27373,5 µl/Test,orange line) or Biotin Rabbit IgG isotype control (A25626,5 µl/Test,blue line), followed by APC Streptavidin (AT0008,0.25 µg) staining. Non-fluorescently stained cells were used as blank control (red line).