

CASP3 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM01771

Basic Information

Catalog No.

RM01771

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

CASP3

Species

Human

Gene ID

836

Swiss Prot

P42574

Synonyms

CPP32; CPP32B; SCA-1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein. [provided by RefSeq, Jul 2008]

Product Information

Description

CASP3 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:46bp deletion in exon2

Allele-2:46bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT AGCGAATCAATGGA*****ATGGGTTTATGTAT
Mut AGCGAATCAATGGA***Deletion***ATGGGTTTATGTAT
Allele-1: 46bp deletion in exon2
WT AGCGAATCAATGGA*****ATGGGTTTATGTAT
Mut AGCGAATCAATGGA***Deletion***ATGGGTTTATGTAT
Allele-2: 46bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CASP3 knockout (KO) 293T cells, using sanger sequencing.