# ABclonal www.abclonal.com

# TET3 Knockout 293T Cell Line, Homozygous

Catalog No.: RM01887

# **Basic Information**

#### Catalog No.

RM01887

# Category

Cell Line

### **Parental Cell line**

293T

# Genotype

Knockout

# **Background**

Members of the ten-eleven translocation (TET) gene family, including TET3, play a role in the DNA methylation process (Langemeijer et al., 2009 [PubMed 19923888]).[supplied by OMIM, Nov 2010]

# **Gene Information**

# **Gene Symbol**

TET3

#### **Species**

Human

# Gene ID

200424

# **Swiss Prot**

043151

# **Synonyms**

hCG 40738

# **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Product Information**

#### Description

TET3 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:194bp deletion in exon3

Allele-2:196bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

# **Packaging**

 ${\bf 1}$  vial parental cell line and  ${\bf 1}$  vial knockout cell line

# **Shipping Conditions**

**Amount** 

Dry ice

1~5x10<sup>6</sup> cells/vial

## Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

#### Protoco

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at  $37^{\circ}C$  with 5% CO<sub>2</sub> condition.

- 1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

# Sequencing data

WT GGAAGATGCCCACG\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*CGGCATGGTATGAA
Mut GGAAGATGCCCACG\*\*\*Deletion\*\*\*\*CGGCATGGTATGAA
Allele-1: 194bp deletion in exon3

WT CCTGGAAGATGCCC\*\*\*\*\*\*\*\*\*\*GCGGCATGGTATGA
Mut CCTGGAAGATGCCC\*\*\*Deletion\*\*\*GCGGCATGGTATGA

Allele-2: 196bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and TET3 knockout (KO) 293T cells, using sanger sequencing.