

LDHA Knockdown 293T Cell Line, Heterozygous

Catalog No.: RM01890

Basic Information

Catalog No.

RM01890

Category

Cell Line

Parental Cell line

293T

Genotype

Knockdown

Gene Information

Gene Symbol

LDHA

Species

Human

Gene ID

3939

Swiss Prot

P00338

Synonyms

GSD11; HEL-S-133P; LDHM; PIG19

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The protein encoded by this gene catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of anaerobic glycolysis. The protein is found predominantly in muscle tissue and belongs to the lactate dehydrogenase family. Mutations in this gene have been linked to exertional myoglobinuria. Multiple transcript variants encoding different isoforms have been found for this gene. The human genome contains several non-transcribed pseudogenes of this gene. [provided by RefSeq, Sep 2008]

Product Information

Description

LDHA Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:exon2 was deleted

Allele-2:WT

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount1~5x10⁶ cells/vial**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT TTCTAAAAATCTAG*****CTTAGGGTAGAGTG
Mut TTCTAAAAATCTAG***Deletion***CTTAGGGTAGAGTG
Allele-1: exon2 was deleted
WT GAAGTGGCAATTTCCATTAACTAAAGATTGATGTC
Mut GAAGTGGCAATTTCCATTAACTAAAGATTGATGTC
Allele-2: WT

Genome sequence analysis of PCR products from parental (WT) and LDHA Knockdown (KD) 293T cells, using sanger sequencing.