

# MDM2 Knockdown HCT116 Cell Line, Heterozygous

Catalog No.: RM01901

## Basic Information

**Catalog No.**

RM01901

**Category**

Cell Line

**Parental Cell line**

HCT116

**Genotype**

Knockdown

## Gene Information

**Gene Symbol**

MDM2

**Species**

Human

**Gene ID**

4193

**Swiss Prot**

Q00987

**Synonyms**

ACTFS; HDMX; hdm2

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes a nuclear-localized E3 ubiquitin ligase. The encoded protein can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. This gene is itself transcriptionally-regulated by p53. Overexpression or amplification of this locus is detected in a variety of different cancers. There is a pseudogene for this gene on chromosome 2. Alternative splicing results in a multitude of transcript variants, many of which may be expressed only in tumor cells. [provided by RefSeq, Jun 2013]

## Product Information

**Description**

MDM2 Knockdown HCT116 Cell Line is engineered from HCT116 cell line with Gene-Editing Technology.

Allele-1:24bp deletion in exon1

Allele-2:38bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**

Dry ice

**Amount**

1~5x10<sup>6</sup> cells/vial

**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

**Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT ATGCTGTACCTAC\*\*\*\*\*GATTCCAGCTTCGG  
Mut ATGCTGTACCTAC\*\*\*Deletion\*\*\*GATTCCAGCTTCGG  
Allele-1: 24bp deletion in exon1  
WT AATACCAACATGTC\*\*\*\*\*CAGCTTCGGAACAA  
Mut AATACCAACATGTC\*\*\*Deletion\*\*\*CAGCTTCGGAACAA  
Allele-2: 38bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and MDM2 Knockdown (KD) HCT116 cells, using sanger sequencing.