

TEAD1 Knockout HeLa Cell Line, Homozygous

Catalog No.: RM01923

Basic Information

Catalog No.

RM01923

Category

Cell Line

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

TEAD1

Species

Human

Gene ID

7003

Swiss Prot

P28347

Synonyms

AA; NTEF-1; REF1; TCF-13; TCF13;
TEAD-1; TEF-1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene encodes a ubiquitous transcriptional enhancer factor that is a member of the TEA/ATTS domain family. This protein directs the transactivation of a wide variety of genes and, in placental cells, also acts as a transcriptional repressor. Mutations in this gene cause Sveinsson's chorioretinal atrophy. Additional transcript variants have been described but their full-length natures have not been experimentally verified. [provided by RefSeq, May 2010]

Product Information

Description

TEAD1 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:23bp insertion and 118bp deletion in exon1

Allele-2:70bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT TGACTCTG*****GAAGGCAA
Mut TGACTCTG***Insertion*****Deletion***GAAGGCAA
Allele-1: 23bp insertion and 118bp deletion in exon1

WT TCTGGAGCCCGAC*****ACGAAGGCAAAATG
Mut TCTGGAGCCCGAC***Deletion***ACGAAGGCAAAATG
Allele-2: 70bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and TEAD1 knockout (KO) HeLa cells, using sanger sequencing.