

# NLRP3 Knockout 293T Cell Line, Homozygous

Catalog No.: RM01936

## Basic Information

### Catalog No.

RM01936

### Category

Cell Line

### Parental Cell line

293T

### Genotype

Knockout

## Gene Information

### Gene Symbol

NLRP3

### Species

Human

### Gene ID

114548

### Swiss Prot

Q96P20

### Synonyms

AGTAVPRL; All; AVP; C1orf7; CIAS1;  
CLR1.1; FCAS; FCAS1; FCU; MWS; NALP3;  
PYPAF1

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and neonatal-onset multisystem inflammatory disease (NOMID). Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Alternative 5' UTR structures are suggested by available data; however, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid. [provided by RefSeq, Oct 2008]

## Product Information

### Description

NLRP3 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:64bp deletion in exon1

Allele-2:85bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### Packaging

1 vial parental cell line and 1 vial knockout cell line

### Shipping Conditions

Dry ice

### Amount

1~5x10<sup>6</sup> cells/vial

### Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

### Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT GATCTAGCCACGCT\*\*\*\*\*AACAGGAGAGACCT  
Mut GATCTAGCCACGCT\*\*\*Deletion\*\*\*AACAGGAGAGACCT  
Allele-1: 64bp deletion in exon1  
WT TGGATCTAGCCACG\*\*\*\*\*AGAAAGCAAAAAGA  
Mut TGGATCTAGCCACG\*\*\*Deletion\*\*\*AGAAAGCAAAAAGA  
Allele-2: 85bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and NLRP3 knockout (KO) 293T cells, using sanger sequencing.