Leader in Biomolecular Solutions for Life Science

www.abclonal.com

SMAD9 Knockout 293T Cell Line, Homozygous

Catalog No.: RM01956

Basic Information

Catalog No.
RM01956

Category
Cell Line

Parental Cell line
293T

Genotype
Knockout

Background

The protein encoded by this gene is a member of the SMAD family, which transduces signals from TGF-beta family members. The encoded protein is activated by bone morphogenetic proteins and interacts with SMAD4. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jan 2010]

Gene Information

Gene Symbol
SMAD9

Species
Human

Gene ID
4093

Swiss Prot
O15198

Synonyms
MADH6; MADH9; PPH2; SMAD8; SMAD8/9; SMAD8A; SMAD8B

Product Information

Description
SMAD9 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.
Allele-1:82bp deletion in exon1
Allele-2:82bp deletion in exon1
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial

Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.


Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and SMAD9 knockout (KO) 293T cells, using sanger sequencing.