

DCN Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM01996

Basic Information

Catalog No.

RM01996

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

DCN

Species

Human

Gene ID

1634

Swiss Prot

P07585

Synonyms

CSCD; DSPG2; PG40; PGII; PGS2; SLRR1B

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene encodes a member of the small leucine-rich proteoglycan family of proteins. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature protein. This protein plays a role in collagen fibril assembly. Binding of this protein to multiple cell surface receptors mediates its role in tumor suppression, including a stimulatory effect on autophagy and inflammation and an inhibitory effect on angiogenesis and tumorigenesis. This gene and the related gene biglycan are thought to be the result of a gene duplication. Mutations in this gene are associated with congenital stromal corneal dystrophy in human patients. [provided by RefSeq, Nov 2015]

Product Information

Description

DCN Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:77bp deletion in exon1

Allele-2:77bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CCTGGGCTGGACCG*****CGACTTCGAGCCCT
Mut CCTGGGCTGGACCG***Deletion***CGACTTCGAGCCCT
Allele-1: 77bp deletion in exon1

WT CCTGGGCTGGACCG*****CGACTTCGAGCCCT
Mut CCTGGGCTGGACCG***Deletion***CGACTTCGAGCCCT
Allele-2: 77bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and DCN knockout (KO) HeLa cells, using sanger sequencing.