

CYP11A1 Knockdown HeLa Cell Lysate, Heterozygous

Catalog No.: RM01997

Basic Information

Catalog No.

RM01997

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockdown

Gene Information

Gene Symbol

CYP11A1

Species

Human

Gene ID

1583

Swiss Prot

P05108

Synonyms

CYP11A; CYPXIA1; P450SCC

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the mitochondrial inner membrane and catalyzes the conversion of cholesterol to pregnenolone, the first and rate-limiting step in the synthesis of the steroid hormones. Two transcript variants encoding different isoforms have been found for this gene. The cellular location of the smaller isoform is unclear since it lacks the mitochondrial-targeting transit peptide. [provided by RefSeq, Jul 2008]

Product Information

Description

CYP11A1 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:69bp deletion in exon2

Allele-2:70bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CCGAGGCCAGCG*****CCGTCTGTTAGGA
Mut CCGAGGCCAGCG***Deletion***CCGTCTGTTAGGA
Allele-1: 69bp deletion in exon2

WT CCGAGGCCAGCG*****CGTCTGTTAGGAC
Mut CCGAGGCCAGCG***Deletion***CGTCTGTTAGGAC
Allele-2: 70bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CYP11A1 Knockdown (KD) HeLa cells, using sanger sequencing.