

# FOS Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02011

## Basic Information

**Catalog No.**

RM02011

**Category**

Cell Lysate

**Parental Cell line**

HeLa

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

FOS

**Species**

Human

**Gene ID**

2353

**Swiss Prot**

P01100

**Synonyms**

AP-1; C-FOS; p55

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death. [provided by RefSeq, Jul 2008]

## Product Information

**Description**

FOS Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.  
Allele-1:139bp deletion in exon2  
Allele-2:140bp deletion in exon2  
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.  
Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.  
Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**

4°C

**Amount**

50µL, 2µg/µL.

**Storage**

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

**Protocol**

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT GGTCACTGCCATCT\*\*\*\*\*CATGACAGGAGGCC  
Mut GGTCACTGCCATCT\*\*\*Deletion\*\*\*CATGACAGGAGGCC  
Allele-1: 139bp deletion in exon2  
WT GGTCACTGCCATCT\*\*\*\*\*ATGACAGGAGGCCG  
Mut GGTCACTGCCATCT\*\*\*Deletion\*\*\*ATGACAGGAGGCCG  
Allele-2: 140bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and FOS knockout (KO) HeLa cells, using sanger sequencing.