XBP1 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM02023



Basic Information

Catalog No. RM02023

Category Cell Lysate

Parental Cell line 293T

Genotype Knockout

Gene Information

Gene Symbol XBP1

Species Human

Gene ID 7494

Swiss Prot P17861

Synonyms TREB-5; TREB5; XBP-1; XBP2

Contact

| 6 | 400-999-6126 |
|----------|---------------------------|
| \times | cn.market@abclonal.com.cn |
| € | www.abclonal.com.cn |

Background

This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor. The isoform encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of XBP1 has been identified and localized to chromosome 5. [provided by RefSeq, Jul 2008]

Product Information

Description

XBP1 Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:exon2 was deleted

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

| Shipping | Conditions |
|----------|------------|
| 4℃ | |

Amount 50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in $1 \times$ SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT TTTGATATGACTCA***********TATACCCTTCCTTG Mut TTTGATATGACTCA***Deletion***TATACCCTTCCTTG Allele-1: exon2 was deleted

WT TTTGATATGACTCA***********TATACCCTTCCTTG Mut TTTGATATGACTCA***Deletion***TATACCCTTCCTTG Allele-2: exon2 was deleted Genome sequence analysis of PCR products from parental (WT) and XBP1 knockout (KO) 293T cells, using sanger sequencing.