

# IL6R Knockout HCT116 Cell Lysate, Homozygous

**Catalog No.:** RM02026

## Basic Information

**Catalog No.**

RM02026

**Category**

Cell Lysate

**Parental Cell line**

HCT116

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

IL6R

**Species**

Human

**Gene ID**

3570

**Swiss Prot**

P08887

**Synonyms**CD126; IL-6R-1; IL-6RA; IL6Q; IL6RA;  
IL6RQ; gp80

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes a subunit of the interleukin 6 (IL6) receptor complex. Interleukin 6 is a potent pleiotropic cytokine that regulates cell growth and differentiation and plays an important role in the immune response. The IL6 receptor is a protein complex consisting of this protein and interleukin 6 signal transducer (IL6ST/GP130/IL6-beta), a receptor subunit also shared by many other cytokines. Dysregulated production of IL6 and this receptor are implicated in the pathogenesis of many diseases, such as multiple myeloma, autoimmune diseases and prostate cancer. Alternatively spliced transcript variants encoding distinct isoforms have been reported. A pseudogene of this gene is found on chromosome 9.[provided by RefSeq, May 2011]

## Product Information

**Description**

IL6R Knockout HCT116 Cell Line is engineered from HCT116 cell line with Gene-Editing technology.

Allele-1:151bp deletion in exon2

Allele-2:151bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**

4°C

**Amount**

50µL, 2µg/µL.

**Storage**

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

**Protocol**

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT GACTCTGACCTGCC\*\*\*\*\*GGCCGGCCGCCAG  
Mut GACTCTGACCTGCC\*\*\*Deletion\*\*\*GGCCGGCCGCCAG  
Allele-1: 151bp deletion in exon2  
WT GACTCTGACCTGCC\*\*\*\*\*GGCCGGCCGCCAG  
Mut GACTCTGACCTGCC\*\*\*Deletion\*\*\*GGCCGGCCGCCAG  
Allele-2: 151bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and IL6R knockout (KO) HCT116 cells, using sanger sequencing.