

FGF2 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02047

Basic Information

Catalog No.

RM02047

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

FGF2

Species

Human

Gene ID

2247

Swiss Prot

P09038

Synonyms

BFGF; FGF-2; FGFB; HBGF-2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF. [provided by RefSeq, Jul 2008]

Product Information

Description

FGF2 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:4bp deletion in exon1

Allele-2:71bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CACCACGCTGCCCCGCTTGCCCGAGGATGGCGGCAGCG
Mut CACCACGCTGCCCCGCT- - -CGAGGATGGCGGCAGCG
Allele-1: 4bp deletion in exon1
WT CCCCgcgcggctcc*****TGCCCGAGGATGGC
Mut CCCCgcgcggctcc***Deletion***TGCCCGAGGATGGC
Allele-2: 71bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and FGF2 Knockout (KO) HeLa cells, using sanger sequencing.